



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

gether by detached pairs, although in some places where unusually abundant they gather daily for a sociable feed. — WILLIAM L. DAWSON, *Oberlin, Ohio.*

A New Bird for the Virginias.— I have lately received from Mr. Thaddeus Surber, White Sulphur Springs, West Virginia, a young male specimen of the Stilt Sandpiper (*Micropalama himantopus*) which was taken on November 2, 1896, near Caldwell, P. O., on the Greenbrier River, six miles from the above mentioned Springs. This species has not hitherto been reported from either Virginia or West Virginia, although a bird from the Patuxent River, Md., has been recorded by Mr. Hugh M. Smith (Auk, Vol. III, p. 139). — WILLIAM C. RIVES, M. D., *New York City.*

Asarcia spinosa.— In Vol. XXIV of the British Museum Catalogue of Birds, Limicolæ, p. 86, 1896, Dr. Sharpe names a genus *Asarcia*, type *Parra variabilis* Linn., 1766, as distinguished from *Jacana* proper by the reduction of the facial appendages to a comparatively small frontal leaf, which is trifid, somewhat like a fleur-de-lis; the same cutaneous excrescence in the type of *Jacana* being bifid or heart-shaped, and supplemented by a pair of wattles. Assuming the generic sufficiency of this distinction, I may remark that Dr. Sharpe's use of the specific term *variabilis* in this connection is at variance with A. O. U. canons of nomenclature. As shown by Elliot, Auk, July, 1888, p. 298, and as admitted by Sharpe, *I. c.*, *Parra variabilis* Linn., S. N. 1766, p. 260, was based on Edwards's Nat. Hist. I. 1743, p. and pl. 48, as a mere renaming of *Fulica spinosa* Linn. S. N. 1758, p. 152, which had the identical basis of Edwards's pl. 48. Consequently *variabilis* is untenable by our rules, and the species should continue to stand as *Jacana spinosa*, unless we recognize the new generic name; in which probable contingency *Asarcia spinosa* becomes the onym of the bird. — ELLIOTT COUES, *Washington, D. C.*

The Passenger Pigeon (*Ectopistes migratorius*) in Lewis County, N. Y.— In the Boonville (N. Y.) 'Herald' of May 27, 1896, appeared an item to the effect that Mr. Henry Felshaw had recently seen "a large flock of Wild Pigeons, perhaps 300 in number, flying westward." On writing to Mr. Felshaw he replied that there was no possibility of his having been mistaken, as he had shot, trapped and netted thousands of these birds, in former years, and the flock in question was seen when not more than fifteen rods distant. The flock, as said, numbered about 300, and was seen at about 5.30 A. M. on May 22, the locality being Constableville, Lewis County, N. Y. He further states, "I mounted, last spring (in April, I think) a cock Pigeon that was shot somewhere near North Western." — W. S. JOHNSON, *Boonville, N. Y.*

Melopelia leucoptera in Osceola County, Florida.— The National Museum has recently received a wing and foot of this species from Prof. W.